


Health monitoring

Guide for organophosphate pesticides





Contents

Orga	anophosphate pesticides	5
1. Sa	Health monitoring for organophosphate pesticides under the Work Health and afety (WHS) Regulations	6
2.	Monitoring exposure to OPs	6
(Other health monitoring methods	8
١	Workplace exposure standard	9
ı	Removal from work	9
ı	Return to work	9
3.	Final medical examination	. 10
4.	Route of occupational exposure	. 10
5.	Target organ/effect	. 11
6.	Acute effects	. 11
7.	Chronic effects	. 14
8.	Carcinogenicity	. 14
9.	GHS classifications	. 14
So	ource documents	. 14
Heal	Ith monitoring report – Organophosphate pesticides	. 17
Se	ection 1 – A copy of this section to be provided to the PCBU	. 17
Se	ection 2 – This section to be retained by the registered medical practitioner	. 21

Introduction

This guide is intended to be read by a registered medical practitioner with experience in health monitoring who is engaged by person conducting a business or undertaking (PCBU) to carry out or supervise health monitoring. It provides practical guidance to registered medical practitioners about requirements under the work health and safety (WHS) laws for health monitoring.

This guide applies to all workplaces covered by the WHS Regulations where health monitoring is required.

How to use this guide

This guide includes references to the legal requirements under the WHS Act and WHS Regulations. These are included for convenience only and should not be relied on in place of the full text of the WHS Act or WHS Regulations.

The words 'must', 'requires' or 'mandatory' indicate a legal requirement exists that must be complied with. The word 'should' is used in this guide to indicate a recommended course of action, while 'may' is used to indicate an optional course of action.

This guide provides information for those registered medical practitioners engaged by a PCBU to carry out or supervise health monitoring for workers. This guidance should be read in conjunction with the following:

- Health monitoring guide for registered medical practitioners
- Health monitoring guides for hazardous chemicals
- Health monitoring guide for workers
- Health monitoring guide for persons conducting business or undertakings (PCBUs).

Health monitoring under the WHS Regulations

In certain circumstances, the model WHS Regulations place duties on a PCBU to provide health monitoring to workers. These requirements arise if the worker is carrying out work with hazardous chemicals including lead and asbestos. In addition, the work being carried out must be the kind of work specified in the WHS Regulations. A PCBU has the duty to determine if health monitoring is required.

The WHS Regulations prescribe that health monitoring is carried out by or supervised by a registered medical practitioner with experience in health monitoring.

Organophosphate pesticides

Organophosphate pesticides are phosphotriester compounds that can be divided into five classes based on their differing alkyl phosphate constituents:

- phosphates
 - chlorfenvinphos
 - o paraoxon
- O-phosphorothioates
 - o chlorpyrifos
 - o fenthion
 - parathion
- S-phosphorothioates
 - Azamethiphos
 - o omethoate
- phosphorodithioates
 - ethion
 - malathion
 - o phosmet, and
- phosphonates/phosphoramidates
 - o acephate, and
 - o fenamiphos.

Work activities that may represent a high risk exposure

Most organophosphorus compounds are insecticides, though there are also a number of related herbicide and fungicide compounds. Organophosphate insecticides are widely used on a large variety of crops.

Examples of work activities involving OPs that require special attention when assessing exposure include:

- pest control operators who use OPs every day in their work
- manufacture, formulation and packaging
- transport, storage and distribution
- handling used containers, for example in scrap recovery
- agricultural and horticultural activities like mixing, loading and applications where direct handling of the chemical occurs
- veterinary activities like cattle and sheep dipping
- seasonal field workers exposed to pesticide residues, and
- laboratory workers undertaking research on OPs.

To be legally used in Australia OP products must be registered by the Australian Pesticides and Veterinary Medicines Authority (APVMA). Product registrations will change over time and referring to the product label and safety data sheet to determine whether the chemical products used contain cholinesterase inhibiting (anti-cholinesterase) compounds is recommended.

Sources of non-occupational exposure

Many OPs are commonly used in home and garden products. For example, diazinon and chlorpyrifos have been widely used by consumers who may contaminate themselves or their food by not understanding or taking the precautions necessary for safe use.

Cases have been reported of acetylcholinesterase inhibition by:

- alkyl sulfates
- sulfonates, and

- a wide variety of drugs for example:
 - o neostigmine
 - o physostigmine
 - o pyridostigmine
 - o pethidine
 - o some immunosuppressants. and
 - various cytostatic agents.

1. Health monitoring for organophosphate pesticides under the Work Health and Safety (WHS) Regulations

Collection of demographic, medical and occupational history, including pattern of use Physical examination

Urinary organophosphate metabolites

Health monitoring before starting work with organophosphate pesticides

Health monitoring for organophosphate pesticides (OPs) may be required before the worker starts work so that changes to the worker's health can be detected.

Initial discussions about a health monitoring program should include:

- possible health effects from exposure to OPs
- how to recognise and report symptoms, and
- what is involved in the health monitoring program, for example the frequency of testing and the tests that may be needed.

A physical examination should be carried out only if work and medical history indicates this is necessary, for example by the presence of relevant symptoms. OPs are respiratory irritants, therefore it is important to investigate respiratory symptoms. However, spirometry may not be required at this stage.

If erythrocyte or plasma cholinesterase activity levels are monitored during the health monitoring program, it is recommended baseline levels are taken. As there is considerable inter-individual variability in baseline cholinesterase activity, an individual baseline must be established for comparison purposes. If the worker has had previous exposure, it is desirable that there is a two month exposure-free period before baseline samples are taken. Two blood samples should be collected at least three days apart before exposure begins. The difference between samples should be less than 20 per cent.

During exposure to an organophosphate pesticide process

2. Monitoring exposure to OPs

Where workers are exposed, suspected of being exposed or are concerned about exposure to OPs, the person conducting the business or undertaking (PCBU) has a duty to arrange a health monitoring appointment with a registered medical practitioner. For example, an appointment should be arranged following spills or loss of containment of OPs resulting in excessive exposure to workers or when workers develop symptoms of OP exposure.

OPs are readily absorbed by inhalation, through the skin and by ingestion. The dermal route is thought to be the major route of occupational exposure for most OPs. The rate and extent of absorption is dependent on the OP and the formulation.

OPs undergo two generalised metabolic reactions:

- hydrolysis to an alkyl phosphate and another compound that may subsequently undergo further metabolism, and
- oxidation of P=S groups in thion OPs (O-phosphorothioates and phosphorodithioates) to P=O groups.

There are six dialkyl phosphate compounds that, collectively, are metabolites of most OPs:

- dimethylphosphate (DMP)
- dimethylthiophosphate (DMTP)
- dimethyldithiophosphate (DMDTP)
- diethylphosphate (DEP)
- diethylthiophosphate (DETP), and
- diethyldithiophosphate (DEDTP).

The following table highlights the potential metabolites from common OPs:

Table 1 Possible dialkyl metabolites from common organophosphate pesticides¹

Metabolite(s)*	Organophosphate pesticide	
DEP DETP	Chlorfenvinphos Chlorpyrifos Diazinon	Parathion pirimiphos-methyl Pyrazophos
DEP DETP DEDTP	Azinphos-ethyl Ethion	Phorate Terbufos
DMP	Dichlorvos Mevinphos	Monocrotophos Trichlophon
DMP DMTP	Azamethiphos Chlorpyrifos-methyl Famphur Fenitrothio Fenthion	Omethoate Parathion-methyl Temephos Toclfos-methyl Vamidothion
DMP DMTP DMDTP	Azinphos-methyl Dimethoate Malathion	Methidathion Phosmet

^{*}One or more of these metabolites would be expected

The metabolites of OPs are excreted into urine or into faeces through biliary excretion.

The following test should be used to test the worker's OP exposure levels:

urinary organophosphate metabolite (dialkyl phosphate) levels.

No biological exposure guidance values have been published. The following values may be used as a guide for interpreting results:

Biological exposure guide for organophosphate pesticides¹

Urinary dialkyl phosphate: less than 100 µmol/mol creatinine

indicates low occupational exposure or high non-occupational exposure

Urinary dialkyl phosphate: 100-1000 µmol/mol creatinine — indicates **medium level occupational exposure**

Urinary dialkyl phosphate: greater than 1000 µmol/mol creatinine

- may be associated with a drop in the blood cholinesterase level
- indicates high occupational exposure

Urinary dialkyl phosphate levels greater than 100 µmol/mol creatinine indicate that work practices may need to be reviewed. An assessment of erythrocyte or plasma cholinesterase activity may be warranted.

While the pharmacokinetic profile of most OPs is not known, it is recommended that urine samples are collected post-shift or pre-shift the following day.

Other health monitoring methods

OPs inhibit acetylcholinesterase (AChE) in the synapses of the nervous system. This is the primary mechanism for the prominent toxicity effects associated with these compounds. Inhibition of the synaptic AChE enzyme is paralleled by inhibition of erythrocyte AChE and plasma cholinesterase. Monitoring the activity of the latter two enzymes can provide an indication of early signs of OP over-exposure. A reduction in cholinesterase activity correlates with clinical symptoms, though there is a wide safety margin between a detectable change in cholinesterase level and the development of significant symptoms.

The cholinesterase test is not as sensitive as the urinary metabolite test. While low plasma or erythrocyte activity indicates OP exposure, no firm conclusions can be made when normal levels of activity for these enzymes are observed.

Due to the considerable differences in baseline activities observed between individuals (potentially greater than 30 per cent difference), baseline activities should be obtained for each worker. Activities of erythrocyte AChE and plasma cholinesterase obtained during the exposure period are then compared with an individual's baseline activity level. The tests should be performed by the same laboratory and same method as that used to obtain the baseline activities.

If erythrocyte AChE activity is monitored, the following value should be used as a guide:

Acetylcholinesterase activity in erythrocytes²:

70 per cent of individual's baseline activity

Cholinergic health effects may be seen at lower AChE activities. Sampling should be performed as soon as possible after exposure or at the end of shift if this is also the end of the exposure period.

The kinetics of erythrocyte AChE and plasma cholinesterase inhibition differ depending on the OP and are different between the two enzymes. Thion OPs require activation to oxon OPs (oxidation of P=S groups to P=O groups) before inhibition of the cholinesterase enzymes can occur. Therefore, a delay in inhibition of erythrocyte AChE and plasma

¹ See <u>Chemical analysis branch handbook, 9th Edition, Workplace and biological monitoring exposure analysis, WorkCover NSW (PDF 3.39MB) for more details</u>

² American Conference of Governmental Industrial Hygienists (ACGIH) (2017) Documentation of the Biological Exposure Indices for Chemical Agents, Acetylcholinesterase Inhibiting Pesticides, 7th Ed, Cincinnati.

cholinesterase activity may be seen with O-phosphorothioate and phosphorodithioate OPs compared with the other classes of OPs.

Cholinesterase inhibition by OPs can be long lasting and can provide an indication of chronic exposure as well as high acute exposures. Following inhibition, a small set of both erythrocyte AChE and plasma cholinesterase undergo spontaneous reactivation, but the majority of the inhibited enzymes 'age' to inactive enzymes. For an aged enzyme, activity can only be restored by replacement with a new enzyme. The rates of reactivation and aging differ between each of the cholinesterase enzymes and the inhibiting OP. Aged plasma cholinesterase is replaced with a half-life of six to 12 days, while aged erythrocyte AChE is replaced with a half-life of approximately 30 days. This long inhibition needs to be considered when resampling to ascertain recovery of enzyme activity.

The frequency of ongoing monitoring should be determined based on an assessment of the type of work, pattern of exposure, the toxicity of the pesticide(s) being handled or used and work practices. Use patterns and potential incidents of exposure should be recorded. If a significant depression of erythrocyte or plasma cholinesterase activity is observed, the individual should be monitored for signs and symptoms of OP exposure, with particular attention to the nervous system.

Rural communities may have some difficulties with collection, transport and storage of blood samples. An arrangement or plan could be negotiated with a local hospital or authorised doctor, so that blood tests suitable for monitoring OP exposure are carried out at a particular seasonal time and then at an appropriate time of the day or week, for occasional or intermittent users. This arrangement would cover the majority of situations.

There is a small proportion of the population that has low plasma cholinesterase activity for genetic reasons. Also, some medical conditions can affect erythrocyte AChE or plasma cholinesterase activity. These factors should be considered when interpreting results.

Monitoring changes in cholinesterase activity is a non-specific means of assessing OP exposure. There are other pesticides, such as carbamates and pyrethroids, and non-pesticides, such as some drugs, that may also inhibit cholinesterase activity.

Workplace exposure standard

Different OPs have different workplace exposure standards.

A physical examination may be indicated if the results of air monitoring indicate frequent or potentially high exposure (half of the eight hour time weighted average (TWA) or above).

Removal from work

Where a medical examination indicates the worker is displaying symptoms of exposure to OPs or where results of biological monitoring indicate exposure that may cause adverse health effects, the registered medical practitioner should consider recommending the worker be removed from OP-related work.

When removal from OP-related work is indicated the registered medical practitioner must provide the PCBU with the following recommendations:

- the worker should be removed from work with OPs the worker can be moved to another area or can use other classes of pesticides, except pyrethroids (such as permethrin) and carbamates, and
- the PCBU should review control measures and carry out recommended remedial action.

The worker must be informed of the results of health monitoring.

Return to work

Should a worker be removed from OP-related work, they must not return until the registered medical practitioner has:

- assessed them as medically fit, and
- made a recommendation to the PCBU that the worker can return to remediated OP-related work.

This assessment should take into consideration the clinical condition of the worker, the worker's urinary dialkyl phosphate levels, erythrocyte or plasma cholinesterase activity and remediation of the circumstances that led to the symptoms if possible.

At termination of work in an organophosphate pesticide process

3. Final medical examination

A urine sample and a blood sample should be collected on the last day of the worker's final shift, and a final medical examination should be carried out at the same time or as soon as possible thereafter. Emphasis should be placed on the nervous system and any symptoms reported during the health monitoring program.

Workers with health conditions or continuing symptoms due to OP exposure should be advised to seek continuing medical examinations as organised by the registered medical practitioner supervising the health monitoring program.

A health monitoring report from the registered medical practitioner should be provided to the PCBU as soon as practicable after the completion of the monitoring program, and at regular intervals for longer term or ongoing health monitoring processes. The report must include:

- the name and date of birth of the worker
- the name and registration number of the registered medical practitioner
- the name and address of the PCBU who commissioned the health monitoring
- the date of the health monitoring
- any test results that indicate whether or not the worker has been exposed to a hazardous chemical
- any advice that test results indicate that the worker may have contracted an injury, illness or disease as a result of carrying out the work that triggered the requirement for health monitoring
- any recommendation that the PCBU take remedial measures, including whether the worker can continue to carry out the type of work that triggered the requirement for health monitoring, and
- whether medical counselling is required for the worker in relation to the work that triggered the requirement for health monitoring.

Potential health effects following exposure to organophosphate pesticides

4. Route of occupational exposure

For most OPs, the dermal route is the primary route of exposure in the workplace.

Inhalation of OPs depends on the volatility of the compound, on the type of formulation and on the technique of application, for example spraying.

Oral exposure may be observed in cases of accidental or deliberate ingestion. Work-related accidental ingestion may occur as a result of poor work practices or lack of personal hygiene.

5. Target organ/effect

Organophosphorus compounds owe their toxic effects to the inhibition of AChE activity in nervous tissue.

AChE, under normal physiological conditions, performs the breakdown of acetylcholine; the chemical mediator responsible for physiological transmission of nerve impulses. In the presence of an OP, AChE is phosphorylated and no longer able to break down acetylcholine into choline and acetic acid. The resulting accumulation of acetylcholine in the parasympathetic nerve synapses (muscarinic-like action), the motor end-plate (nicotine-like action) and in the central nervous system is responsible for all typical symptoms occurring after acute poisoning with OP. The synapses initially become overstimulated, before becoming exhausted, with initial symptoms of over activity, and final symptoms of paralysis.

The target organs and potential effects of OP exposure include:

Table 2 Target organs and potential effects of OP exposure

Target organ	Effect
Central nervous system (via inhibition of acetylcholine receptors)	 Headache Anxiety Restlessness Confusion Slurred speech Convulsions Coma Depression of the respiratory and circulatory centres
Peripheral nervous system (via inhibition of muscarinic and nicotinic receptors)	 Salivation Lacrimation Urination Defaecation Gastric cramps Emesis

6. Acute effects

Typical symptoms of acute exposure to OPs include excessive sweating, slurred speech and blurred vision.

The first symptoms of organophosphate poisoning can occur within minutes of exposure to a concentrate or a highly toxic OP. It is common for symptoms to occur an hour or so after inadvertent skin exposure to a working solution.

The symptoms of OP intoxication can be divided into muscarinic-like and nicotinic-like effects, as well as effects on the central nervous system.

Table 3 Acute effects manifestations

Nervous tissue and receptors affected	Manifestations	
Parasympathetic autonomic (r	nuscarinic receptors) post ganglionic nerve fibres	
Exocrine glands	Increased salivationLacrimation, andPerspiration	
Eyes	 Miosis (pinpoint and non-reactive) ptosis Blurring of vision Conjunctival injection, and 'Bloody-tears' 	
Gastrointestinal tract	 Nausea Vomiting Abdominal tightness Swelling and cramps Diarrhoea Tenesmus, and Faecal incontinence 	
Respiratory tract	 Excessive bronchial secretions Rhinorrhoea Wheezing Oedema Tightness in chest Bronchospasms Bronchoconstriction Cough Bradypnoea, and Dyspnoea 	
Cardiovascular system	Bradycardia, andDecrease in blood pressure	
Bladder	 Urinary frequency, and Incontinence	
Parasympathetic and sympathetic autonomic fibres (nicotinic receptors)		
Cardiovascular system	 Tachycardia Pallor Increase in blood pressure, and 	

Nervous tissue and	Manifestations
receptors affected	

Somatic motor nerve fibres (nicotinic receptors)

	 Muscle fasciculations (eyelids, fine facial muscles) Cramps, Restlessness Generalised motor activity Reaction to acoustic stimuli Tremulousness Diminished tendon reflexes Generalised muscle weakness in peripheral and respiratory muscles Paralysis Flaccid or rigid tone Emotional lability, and Ataxia
--	--

Brain (acetylcholine receptors)

Central nervous system	 Drowsiness Lethargy Fatigue Mental confusion Inability to concentrate Headache Pressure in head Generalised weakness Coma with absence of reflexes Tremors Cheyne-Stokes respiration Dyspnoea Convulsions Depression of respiratory centres, and Cyanosis
------------------------	---

Local effects at the site of exposure may occur without symptoms of systemic absorption:

- a splash in the eye may cause blurred vision due to spasm
- inhalation may cause bronchoconstriction and produce an excess of respiratory tract secretions resulting in feelings of chest tightness and a watery nasal discharge, and
- splashes on the skin may cause localised sweating and fasciculations.

For O-phosphorothioate and phosphorodithioates, symptoms usually reach their maximum severity 24 to 48 hours after onset and usually regress over the next one to six days. The onset of symptoms is much quicker for the remaining classes of OPs.

In the case of massive exposure, death usually occurs within 24 hours.

Another short term effect of organophosphates is the 'intermediate syndrome'. This is characterised by transient muscle weakness of the limbs, neck and respiratory muscles, that begins one to four days after a poisoning incident and may continue for up to several weeks.

Delayed polyneuropathy can occur from inhibition of another nervous tissue esterase called neuropathy target esterase. The interval between acute exposure and the onset of neuropathy may be up to four weeks. Initial symptoms are often sensory and consist of tingling and burning sensations in the hands and feet followed by weakness in the lower limbs and ataxia. In severe cases the upper limbs may be affected. There is no specific treatment for this disorder though physiotherapy may limit the muscle wasting that follows denervation.

7. Chronic effects

Symptoms of chronic poisoning usually do not occur until enzyme activity has been reduced to between 60 to 25 per cent of an individual's baseline.

Chronic low level exposures may lead to cumulative effects. Thus, workers continually exposed may be at high risk even at low level exposures.

Chronic exposure to OPs may cause:

- persistent anorexia
- weakness
- ophthalmological effects
- malaise, or
- certain neurobehavioural effects.

Many OPs cause primary irritant dermatitis; only a few, for example parathion and malathion, are known to cause allergic contact dermatitis.

8. Carcinogenicity

Most OPs have not been classified as carcinogenic according to the Globally Harmonized System of Classification and Labelling of Chemicals (GHS). For further information on specific OPs, refer to Safe Work Australia's Hazardous Chemical Information system or the relevant safety data sheet.

9. GHS classifications

Different OPs may have different health hazard classifications. The specific OP that a worker is exposed to will need to be reviewed to ensure appropriate identification of the health hazards. For the GHS classification of a specific OP, refer to Safe Work Australia's Hazardous Chemical Information System or the relevant safety data sheet for detailed information.

Source documents

Alavanja, M.C.R., Sandler, D.P., Lynch, C.F., Knott, C., Lubin, J.H., Tarone, R., Thomas, K., Dosemeci, M., Barker, J., Hoppin, J.A. and Blair, A. (2005) Cancer Incidence in the Agricultural Health Study. *Scand. J. Work Environ. Health* 31 (suppl 1): 39-45.

American Conference of Governmental Industrial Hygienists (ACGIH) (2017) Documentation of the Biological Exposure Indices for Chemical Agents, Acetylcholinesterase Inhibiting Pesticides, 7th Ed, Cincinnati.

<u>Chemical analysis branch handbook, 9th Edition, Workplace and biological monitoring exposure analysis, WorkCover NSW (PDF 3.39MB).</u>

Coggon, D. (2002) Work with Pesticides and Organophosphate Sheep Dips. *Occ. Med.* 52(8): 467-470.

Dementi, B. (1994) Ocular Effects of Organophosphates: A Historical Perspective of Saku Disease. *J. Appl. Toxicol.* 14(2): 119-129.

Ecobichon, D.J. (1996) Toxic Effects of Pesticides in Klaassen CD (ed), *Casarett and Doull's Toxicology The Basic Science of Poisons*, 5th Ed, pp 643-689, McGraw Hill, New York.

European Centre for Ecotoxicology and Toxicology of Chemicals (1998) Organophosphorous Pesticides and Long-Term Effects on the Nervous system, Technical report No. 75, ECOTOC, Brussels.

Jokanovic, M. and Maksimovic, M. (1997) Abnormal Cholinesterase Activity: Understanding and Interpretation. *Eur. J. Clin. Chem . Clin. Biochem.* 35(1): 11-16.

Health and Safety Executive (2005) *Genetic Variation in Susceptibility to Chronic Effects of Organophosphate Exposure*, Health and Safety Executive Research Report 408.

Health and Safety Laboratory (UK) (2013) Guidance on Laboratory Techniques in Occupational Medicine (13th edition).

Jeyaratnam, J. and Maroni, M. (1994) Organophosphorus Compounds. Toxicol. 91: 15-27.

Kamel, F., Engel, L.S., Gladen, B.C., Hoppin, J.A., Alavanja, M.C. and Sandler, D.P. (2005) Neurologic Symptoms in Licensed Private Pesticide Applicators in the Agricultural Health Study. *Environ. Health Perspect.* 113(7): 877-882.

Lauwerys, R.R. and Hoet, P. (2001) *Industrial Chemical Exposure Guidelines for Biological Monitoring*, 3rd Ed, Lewis Publishers, Boca Raton.

Occupational Safety and Health Service, Dept of Labour New Zealand (2000) A Guideline to Promote Best Practice with Organophosphates. www.osh.dol.govt.nz.

Safe Work Australia (2013); *Workplace Exposure Standards for Airborne Contaminants* (PDF 873KB).

Safe Work Australia; *Hazardous Chemicals Information System*.

Wooller K, ed. *Training Manual for WorkCover Authority Authorised Medical Practitioners*, New South Wales WorkCover Authority, Sydney, 1996.



Health monitoring report

Organophosphate pesticides

Health monitoring report – Organophosphate pesticides

This health monitoring report is a confidential health record and must not be disclosed to another person except in accordance with the Work Health and Safety Regulations or with the consent of the worker.

There are two sections. Complete both sections and all questions as applicable.

Section 1 A copy of this section should be forwarded to the person conducting the business or undertaking (PCBU) who has engaged your services.

Section 2 may contain confidential health information. Information that is required to be given to the PCBU should be summarised in Section 1.

Section 1 - A copy of this section to be provided to the PCBU

Person conducting a business or undertaking

Company/organisation name: Click here to enter text.

Site address: Click here to enter text.

Suburb: Click here to enter text.

Postcode: Click here to enter text.

Site Tel: Click here to enter text.

Site Fax: Click here to enter text.

Contact Name: Click here to enter text.

Other businesses or undertakings engaging the worker

□ N/A

(include a separate section for each PCBU)

Company/organisation name: Click here to enter text.

Site address: Click here to enter text.

Suburb: Click here to enter text.

Postcode: Click here to enter text.

Site Tel: Click here to enter text.

Site Fax: Click here to enter text.

Contact Name: Click here to enter text.

Worker details (tick all relevant boxes)

Surname: Click here to enter text. Given names: Click here to enter text.

Date of birth: Click here to enter a date. Sex: ☐ Male ☐ Female

Address: Click here to enter text.

Suburb: Click here to enter text. **Postcode:** Click here to enter text.

Current job: Click here to enter text.

Tel (H): Click here to enter text. Mob: Click here to enter text.

Date started employment: Click here to enter a date.

Employment in organophosphate pesticides risk work (tick all relevant boxes) (information provided by the PCBU)

Wear gloves	□ Yes □ No
Controls	
Other chemicals the worker may be exposed	to: Click here to enter text.
☐ Other (specify):	
☐ Seasonal field work	☐ Laboratory work
☐ Veterinary work	☐ Farming
☐ Aerial crop spraying	☐ Horticultural industry
\square Transport, storage or distribution	☐ Agricultural industry
☐ Pest control	☐ Manufacture and packaging
Organophosphate pesticide industry/use	
Date of assessment: Click here to enter a date.	
Work environment assessment (tick all rele (information provided by the PCBU)	vant boxes)
☐ Minimal – describe: Click here to enter text.	
☐ Brief – less than 1 hour a day	
☐ Short – 1-5 hours in a day	
☐ Long – 6 or more hours in a day	
Duration of exposure	
\square Seasonal (several days a week for a season)	
☐ Infrequent (1 day or less in a work month)	
☐ Occasional (2-3 days in a work month)	
\square Regular (2-3 days in a work week)	
\Box Frequent (daily; 5 or more days in a work week)
Pattern of exposure	
Risk assessment completed: ☐ Yes ☐ No	
When last used: Click here to enter text.	
Type of organophosphate pesticide(s) used:	Click here to enter text.
Worked with organophosphate pesticides sinc	e: Click here to enter a date.
$\hfill\Box$ Current worker continuing in organophosphate	pesticides work
\square New worker but not new to organophosphate per	esticides work
\square New to organophosphate pesticides work	
Type of organophosphate pesticide used (if known	n; please specify): Click here to enter text.

Controls		
Safety goggles/face shield	☐ Yes	□ No
Respirator use	☐ Yes	□ No
Respirator type text.	Click	here to enter
Local exhaust ventilation	□ Yes	□ No
Overalls/work clothing	□ Yes	□ No
Laundering by employer	☐ Yes	□ No
Wash basins and showers (with hot and cold water)	☐ Yes	□ No
Other please specify		
	•	

Health monitoring results

Biological monitoring results

Include/attach test results that indicate whether or not the worker has been exposed **Urine Alkyl Phosphate Metabolites test** (urinary dialkyl phosphate)

DAP (µmol/mol creatinine)	Timing	Comment
	☐ Pre-shift	
	\square Post-shift	
Click here to enter text.	☐ Next day	Click here to enter text.
	☐ Pre-shift	
	\square Post-shift	
Click here to enter text.	\square Next day	Click here to enter text.
	☐ Pre-shift	
	☐ Post-shift	
Click here to enter text.	\square Next day	Click here to enter text.
	(µmol/mol creatinine) Click here to enter text. Click here to enter text.	☐ Pre-shift ☐ Post-shift ☐ Post-shift ☐ Next day ☐ Pre-shift ☐ Post-shift ☐ Post-shift ☐ Post-shift ☐ Post-shift ☐ Pre-shift ☐ Pre-shift ☐ Post-shift ☐ Post-shift ☐ Post-shift

Note:

- 1. less than 100 μmol/mol creatinine considered low work exposure.
- 2. 100-1000 µmol/mol creatinine indicates work exposure review workplace controls to reduce exposure levels
- 3. greater than 1000 µmol/mol creatinine indicates high work exposure; may be associated with a fall in blood cholinesterase levels.

Comments about health monitoring results (for example any early indications or diagnosis of injury, illness or disease): Click here to enter text.

Recommendations (by registered medical practitioner) (tick all relevant boxes)

Further/additional health monitoring for worker

☐ This is the final health monitoring report				
□ Repeat health assessment in Click here to enter text. month(s) / Click here to enter text. week(s)				
☐ Counselling required				
☐ Medical examination by registered medical practitioner. On Click here to enter a date.				
☐ Referred to Medical Specialist (respiratory/dermatology/other). On Click here to enter a date.				
Recommendations to PCBU				
$\hfill\Box$ The worker is suitable for work with organophos	sphate pesticides			
☐ Review workplace controls				
$\hfill\Box$ The worker should be removed from work with On Click here to enter a date.	organophosphate pesticides.			
$\hfill\Box$ The worker is fit to resume work. On Click here	to enter a date.			
\square Biological monitoring results indicate unaccepta	ıbly high exposure levels			
Specialist's name: Click here to enter text.				
Additional comments or recommendations: Cli	ck here to enter text.			
Registered medical practitioner (responsible	e for supervising health monitoring)			
Name: Click here to enter text.				
Signature:				
Date: Click here to enter a date.				
Tel: Click here to enter text.	Fax: Click here to enter text.			
Registration Number: Click here to enter text.				
Medical Practice: Click here to enter text.				
Address: Click here to enter text.				
Suburb: Click here to enter text. Postcode: Click here to enter text.				

of OP products

This form contains confidential information and must not be disclosed to another person except in accordance with the Work Health and Safety Regulations or with consent of the worker

Section 2 – This section to be retained by the registered medical practitioner

Person conducting a business or undertaking				
Company/organisation name: Click here to enter to Suburb: Click here to enter text. Site Tel: Click here to enter text. Contact Name: Click here to enter	ext.	Post	code: Click here to enter text. Fax: Click here to enter text.	
Other businesses or undertak	kings enga	ging the	worker	□ N/A
Company/organisation name: Cli Site address: Click here to enter to Suburb: Click here to enter text. Site Tel: Click here to enter text. Contact Name: Click here to enter	ext.	Post	code: Click here to enter text. Fax: Click here to enter text.	
Worker details (tick all relevant	boxes)			
Surname: Click here to enter text. Date of birth: Click here to enter a			n names: Click here to enter te	xt.
Sex: Male Female Address: Click here to enter text. Suburb: Click here to enter text. Current job: Click here to enter text. Tel (H): Click here to enter text. Date started employment: Click here to organophosphate pestic	kt. here to enter	Post Mob a date.	code: Click here to enter text. : Click here to enter text.	enter text.
Past employment and exposu	re details	(tick all r	elevant boxes)	
Have you ever worked in any of the following jobs? If you answered 'yes' to any of the questions, please advise if you experienced any symptoms such as cough or wheeze or asthma when working. Comments (all 'yes' answers)				
Pest control or fumigation	□ No	□ Yes	Click here to enter text.	·
Manufacture or packaging of OP products	□ No	☐ Yes	Click here to enter text.	
Transport storage or distribution	□ No	□ Voc	Click here to enter text	

			Comments (all 'yes' answers)
Agriculture	□ No	□ Yes	Click here to enter text.
Horticulture	□ No	□ Yes	Click here to enter text.
Farming	□ No	☐ Yes	Click here to enter text.
Aerial crop spraying	□ No	☐ Yes	Click here to enter text.
Veterinary work	□ No	☐ Yes	Click here to enter text.
Seasonal field work	□ No	☐ Yes	Click here to enter text.
Laboratory work	□ No	☐ Yes	Click here to enter text.
Other (please specify)	□ No	□ Yes	Click here to enter text.
General health questionnaire (tick all rel	levant box	es)
			Comments (all 'yes' answers)
Did you suffer any incapacity lasting two weeks or longer in the last two years	□ No	□ Yes	Click here to enter text.
Have you ever had any operations or accidents or been hospitalised for any reason	□ No	☐ Yes	Click here to enter text.
Are you currently being treated by a doctor or other health professional for any illness or injury	□ No	☐ Yes	Click here to enter text.
Are you currently receiving any medical treatment or taking any medications. Please detail.	□ No	☐ Yes	Click here to enter text.
Do you practice personal hygiene at work, for example nail biting, frequency of hand washing, eating or smoking, clean shaven, shower and change into clean clothes at end of shift	□ No	□ Yes	
Specific health questions (tick	all releva	nt boxes)	
Do you have or have you ever ha	nd:		Comments (all 'yes' answers)
Blurred vision or other vision problems	□ No	□ Yes	Click here to enter text.
Itchy eyes, runny or congested nose	□ No	☐ Yes	Click here to enter text.
Loss of hearing or ringing in the ears	□ No	□ Yes	Click here to enter text.
Chest pains or irregular heartbeats	□ No	□ Yes	Click here to enter text.

Do you have or have you ever ha	ıd:		Comments (all 'yes' answers)
High blood pressure or heart disease (including heart attack, heart surgery, murmurs, angina)	□ No	□ Yes	Click here to enter text.
Shortness of breath on exertion	□ No	☐ Yes	Click here to enter text.
Wheezing, bronchitis or asthma now or in the past	□ No	□ Yes	Click here to enter text.
Allergies, hay fever, or allergic bronchitis	□ No	□ Yes	Click here to enter text.
Severe head or spinal injury resulting in hospitalisation	□ No	□ Yes	Click here to enter text.
Severe headaches or migraines	□ No	☐ Yes	Click here to enter text.
Chronic fatigue or tiredness	□ No	□ Yes	Click here to enter text.
Significant weight loss	□ No	□ Yes	Click here to enter text.
Any neurological condition affecting nerves in your feet or hands, your coordination or balance	□ No	□ Yes	Click here to enter text.
Skin disorders or dermatitis	□ No	☐ Yes	Click here to enter text.
Any previous symptoms associated with organophosphate pesticides	□ No	□ Yes	Click here to enter text.
Any other significant health conditions	□ No	□ Yes	Click here to enter text.
Medications currently taken that r	nay affect	blood tes	t results:
			Comments (all 'yes' answers)
Oral contraceptive pill	□ No	☐ Yes	Click here to enter text.
Lithium	□ No	□ Yes	Click here to enter text.
Prednisone/cortisone	□ No	☐ Yes	Click here to enter text.
Propanolol	□ No	□ Yes	Click here to enter text.
General health assessment (if a	applicable)	
Height: Click here to enter text. cm		Weigl	ht: Click here to enter text. kg
BP: Click here to enter text. / Click here to enter text. mmHg			
Urinalysis			
Blood: □ Normal □ Abnormal			
Protein: Click here to enter text.		Refer	red for further testing

Sugar: Click here to enter text.		□ No □] Yes
Cardiovascular system			Medical comments (for all yes/abnormal)
Blood pressure	□ Normal	☐ Abnormal	Click here to enter text.
Heart rate	☐ Normal	☐ Abnormal	Click here to enter text.
Heart sounds	□ Normal	☐ Abnormal	Click here to enter text.
Murmurs present	□ No	☐ Yes	Click here to enter text.
Evidence of cardiac failure/oedema	□ No	□ Yes	Click here to enter text.
Skin			Click have to output tout
Eczema, dermatitis or allergy	□ No		Click here to enter text.
Skin cancer or other abnormali	-		Click here to enter text.
Evidence of nail biting	□ No		Click here to enter text.
Other	□ No	☐ Yes	Click here to enter text.
		9	

Figure 1 Template of the human body to indicate the location of abnormalities

Biological monitoring results

Include/attach at least the previous two test results (if available)

Date	Tests performed	Recommended action or comment
Click here to enter a date.	Click here to enter text.	Click here to enter text.
Click here to enter a date.	Click here to enter text.	Click here to enter text.
Click here to enter a date.	Click here to enter text.	Click here to enter text.
Click here to enter a date.	Click here to enter text.	Click here to enter text.

Other medical history, family medical history, current medication, comments, tests or recommendations (use separate sheet if necessary)

Click here to enter text.

Registered medical practitioner (responsible for supervising health monitoring)

Name: Click here to enter text.

Signature:

Date: Click here to enter a date.

Tel: Click here to enter text. Fax: Click here to enter text.

Registration Number: Click here to enter text.

Medical Practice: Click here to enter text.

Address: Click here to enter text.

Suburb: Click here to enter text. Postcode: Click here to enter text.